

Leukaemia Section

Short Communication

t(16;21)(p11;q22)

Jean Loup Huret

Genetics, Dept Medical Information, UMR 8125 CNRS, University of Poitiers, CHU Poitiers Hospital, F-86021 Poitiers, France (JLH)

Published in Atlas Database: January 2005

Online updated version : <http://AtlasGeneticsOncology.org/Anomalies/t1621.html>


DOI: 10.4267/2042/38161

This article is an update of: Pérot C. t(16;21)(p11;q22). Atlas Genet Cytogenet Oncol Haematol.1998;2(2):62.

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.

© 2005 Atlas of Genetics and Cytogenetics in Oncology and Haematology

Identity



t(16;21)(p11;q22) G- banding - Courtesy Melanie Zenger and Claudia Haeflrich.

Clinics and pathology

Disease

de novo acute non lymphocytic leukemia (ANLL); to be noted is one case of chronic myelogenous leukemia (CML) -blast crisis.

Phenotype/cell stem origin

ANLL cases: mainly M1, M2, M4, M5a, M5b, or M7 ANLL; may be preceded by a myelodysplastic syndrome (MDS).

Epidemiology

About 40 reported cases, mainly found in young adults; children cases are described; median age is about 30 yrs; balanced sex ratio.

Clinics

Blood data: anemia, thrombocytopenia, mild hyperleucocytosis; with high monocytic cell count at times.

Cytology

Myelocytic and monocytoid features are often present; eosinophils in the bone marrow are sometimes abnormal and/or elevated; erythrophagocytosis may be found.

Prognosis

Seems poor: complete remission may not be achieved; there is high incidence of relapse within a year and a median of survival is about 22 months (cases herein reviewed).

Disease

Ewing tumours

Note

t(16;21)(p11;q22) has been found in rare cases of Ewing tumours, a paediatric neoplasm with small round-cells derived from neural crests cells usually associated with translocations involving EWSR1.

Cytogenetics

Ewing tumours are usually associated with a t(11;22)(q24;q12) with 5' EWSR1 - 3' FLI1 involvement, less often associated with t(21;22)(q22;q12) with 5' EWSR1 - 3' ERG involvement, rarely associated with t(2;22)(q36;q12) (5' EWSR1 - 3' FEV) or with t(17;22)(q21;q12) (5' EWSR1-3' ETV4)

Prognosis

Recent treatments have improved the prognosis of Ewing's tumours. The prognosis is mainly determined by the presence of metastases at the time of diagnosis.

Cytogenetics

Additional anomalies

ANLL cases: found solely in about 60% of cases in at least a subclone; associated with +10, +8, or de(9q)/-9 in about 10% of cases each.

Genes involved and proteins

FUS

Location

16p11

Protein

RNA binding protein; member of the TET family, like EWSR1.

ERG

Location

21q22

Protein

ETS transcription factor.

Result of the chromosomal anomaly

Hybrid gene

Description

5' FUS including exons 1 to 6, 7 or 8 - 3' ERG from exon 7, 8 or 9 to C-term.

Fusion protein

Description

N-term FUS transactivation domain fused to the C-term DNA binding ETS domain of ERG.

Oncogenesis

Seems to act as a transcriptional activator.

References

Mecucci C, Bosly A, Michaux JL, Broekaert-Van Orshoven A, Van den Berghe H. Acute nonlymphoblastic leukemia with

bone marrow eosinophilia and structural anomaly of chromosome 16. *Cancer Genet Cytogenet.* 1985 Aug;17(4):359-63

Minamihisamatsu M, Ishihara T. Translocation (8;21) and its variants in acute nonlymphocytic leukemia. The relative importance of chromosomes 8 and 21 to the genesis of the disease. *Cancer Genet Cytogenet.* 1988 Jul 15;33(2):161-73

Berkowicz M, Rosner E, Resnitzky P, Mamon Z, Ben-Bassat I, Ramot B. Acute nonlymphocytic leukemia with t(16;21) *Cancer Genet Cytogenet.* 1990 Jul 1;47(1):139-40

Sadamori N, Yao E, Tagawa M, Nakamura H, Sasagawa I, Itoyama T, Tokunaga S, Ichimaru M, Nakamura I, Kamei T. 16;21 translocation in acute nonlymphocytic leukemia with abnormal eosinophils: a unique subtype. *Acta Haematol.* 1990;84(4):212-6

Marosi C, Bettelheim P, Geissler K, Lechner K, Köller U, Haas OA, Chott A, Hagemeijer A. Translocation (16;21)(p11;q22) in acute monoblastic leukemia with erythrophagocytosis. *Cancer Genet Cytogenet.* 1991 Jul 1;54(1):61-6

Morgan R, Riske CB, Meloni A, Ries CA, Johnson CH, Lemons RS, Sandberg AA. t(16;21)(p11.2;q22): a recurrent primary rearrangement in ANLL. *Cancer Genet Cytogenet.* 1991 May;53(1):83-90

Ferro MR, Cabello P, Garcia-Sagredo JM, Resino M, San Roman C, Larana JG. t(16;21) in a Ph positive CML. *Cancer Genet Cytogenet.* 1992 Jun;60(2):210-1

Maseki N, Miyoshi H, Shimizu K, Homma C, Ohki M, Sakurai M, Kaneko Y. The 8;21 chromosome translocation in acute myeloid leukemia is always detectable by molecular analysis using AML1. *Blood.* 1993 Mar 15;81(6):1573-9

Nobbs MC, Chan-Lam D, Howell RT, Kitchen C, Copplestone JA. Acute non-lymphocytic leukemia with t(16;21). *Cancer Genet Cytogenet.* 1993 Oct 15;70(2):144-5

Okada K, Takeichi M, Uchida H, Shirota T, Sakai N, Ito H. Translocation (16;21)(p11;q22) in acute nonlymphocytic leukemia. *Cancer Genet Cytogenet.* 1994 Jul 1;75(1):60-3

Panagopoulos I, Aman P, Fioretos T, Höglund M, Johansson B, Mandahl N, Heim S, Behrendtz M, Mitelman F. Fusion of the FUS gene with ERG in acute myeloid leukemia with t(16;21)(p11;q22). *Genes Chromosomes Cancer.* 1994 Dec;11(4):256-62

Satoh K, Miura I, Chubachi A, Ohtani H, Hirokawa M, Niitsu H, Miura AB. [Acute monoblastic leukemia (M5a) with dysmegakaryocytopoiesis associated with t(16;21) (p11;q22)]. *Rinsho Ketsueki.* 1994 Feb;35(2):160-4

Scott AA, Head DR, Kopecky KJ, Appelbaum FR, Theil KS, Grever MR, Chen IM, Whittaker MH, Griffith BB, Licht JD. HLA-DR-, CD33+, CD56+, CD16- myeloid/natural killer cell acute leukemia: a previously unrecognized form of acute leukemia potentially misdiagnosed as French-American-British acute myeloid leukemia-M3. *Blood.* 1994 Jul 1;84(1):244-55

Hiyoshi M, Koh KR, Yamane T, Tatsumi N. Acute non-lymphoblastic leukaemia with t(16;21): case report with a review of the literature. *Clin Lab Haematol.* 1995 Sep;17(3):243-6

Hiyoshi M, Yamane T, Hirai M, Tagawa S, Hattori H, Nakao Y, Yasui Y, Koh KR, Hino M, Tatsumi N. Establishment and characterization of IRTA17 and IRTA21, two novel acute non-lymphocytic leukaemia cell lines with t(16;21) translocation. *Br J Haematol.* 1995 Jun;90(2):417-24

Harigae H, Kobayashi M, Mihara A, Watanabe N. Detection of minimal residual disease in cerebro-spinal fluid of a patient

with acute myelogenous leukemia with t(16;21)(p11;q22) translocation by reverse transcriptase-polymerase chain reaction. *Tohoku J Exp Med.* 1997 Dec;183(4):297-302

Kong XT, Ida K, Ichikawa H, Shimizu K, Ohki M, Maseki N, Kaneko Y, Sako M, Kobayashi Y, Tojou A, Miura I, Kakuda H, Funabiki T, Horibe K, Hamaguchi H, Akiyama Y, Bessho F, Yanagisawa M, Hayashi Y. Consistent detection of TLS/FUS-ERG chimeric transcripts in acute myeloid leukemia with t(16;21)(p11;q22) and identification of a novel transcript. *Blood.* 1997 Aug 1;90(3):1192-9

Tamura S, Takemoto Y, Hashimoto-Tamaoki T, Mimura K, Sugahara Y, Senoh J, Furuyama JI, Kakishita E. Cytogenetic analysis of de novo acute myeloid leukemia with trilineage myelodysplasia in comparison with myelodysplastic syndrome evolving to acute myeloid leukemia. *Int J Oncol.* 1998 Jun;12(6):1259-62

Imashuku S, Hibi S, Kuriyama K, Todo S. Hemophagocytosis by leukemic blasts in a case of acute megakaryoblastic leukemia with t(16;21)(p11;q22). *Int J Hematol.* 1999 Jul;70(1):36-9

Raimondi SC, Chang MN, Ravindranath Y, Behm FG, Gresik MV, Steuber CP, Weinstein HJ, Carroll AJ. Chromosomal abnormalities in 478 children with acute myeloid leukemia: clinical characteristics and treatment outcome in a cooperative pediatric oncology group study-POG 8821. *Blood.* 1999 Dec 1;94(11):3707-16

Shikami M, Miwa H, Nishii K, Takahashi T, Shiku H, Tsutani H, Oka K, Hamaguchi H, Kyo T, Tanaka K, Kamada N, Kita K. Myeloid differentiation antigen and cytokine receptor expression on acute myelocytic leukaemia cells with t(16;21)(p11;q22): frequent expression of CD56 and

interleukin-2 receptor alpha chain. *Br J Haematol.* 1999 Jun;105(3):711-9

Okita H, Umezawa A, Fukuma M, Ando T, Urano F, Sano M, Nakata Y, Mori T, Hata J. Acute myeloid leukemia possessing jumping translocation is related to highly elevated levels of EAT/mcl-1, a Bcl-2 related gene with anti-apoptotic functions. *Leuk Res.* 2000 Jan;24(1):73-7

Dastugue N, Lafage-Pochitaloff M, Pagès MP, Radford I, Bastard C, Talmant P, Mozziconacci MJ, Léonard C, Bilhou-Nabéra C, Cabrol C, Capodano AM, Cornillet-Lefebvre P, Lessard M, Mugneret F, Pérot C, Taviaux S, Fenneteaux O, Duchayne E, Berger R. Cytogenetic profile of childhood and adult megakaryoblastic leukemia (M7): a study of the Groupe Français de Cytogénétique Hématologique (GFCH). *Blood.* 2002 Jul 15;100(2):618-26

Shing DC, McMullan DJ, Roberts P, Smith K, Chin SF, Nicholson J, Tillman RM, Ramani P, Cullinane C, Coleman N. FUS/ERG gene fusions in Ewing's tumors. *Cancer Res.* 2003 Aug 1;63(15):4568-76

Heller A, Loncarevic IF, Glaser M, Gebhart E, Trautmann U, Claussen U, Liehr T. Breakpoint differentiation in chromosomal aberrations of hematological malignancies: Identification of 33 previously unrecorded breakpoints. *Int J Oncol.* 2004 Jan;24(1):127-36

This article should be referenced as such:

Huret JL. t(16;21)(p11;q22). *Atlas Genet Cytogenet Oncol Haematol.* 2005; 9(1):34-36.
